

**Vaccination Guide – Protocols for other countries may differ.**

**Please consult with Dr Donna for clarification**

Age	Vaccine
Birth	BCG
	Hep- B Hepatitis B vaccine.
2 months	Diphtheria- Tetanus-acellular Pertussis-inactivated Polio
	Hib- Haemophilus influenzae type b- Hepatitis B vaccine dose 2 Pneumococcal ( <a href="#">Prevenar</a> )
	Meningococcal C vaccine
	Rota virus(optional)
4 months	Diphtheria- Tetanus-acellular Pertussis-inactivated Polio
	Hib-Hep B- Haemophilus influenza type b- Pneumococcal ( <a href="#">Prevenar</a> ) Meningococcal B vaccine Rota virus(optional)
	Diphtheria- Tetanus-acellular Pertussis-inactivated Polio Hib- Haemophilus influenzae type b- Hepatitis B vaccine. Dose 3
	Meningococcal C vaccine Pneumococcal ( <a href="#">Prevenar</a> ) Rota virus(optional)
10 months	Meningococcal C vaccine
	MMR –measles mumps and Rubella Varicella vaccine Pneumococcal ( <a href="#">Prevenar</a> ) Japanese encephalitis vaccine dose one Hepatitis A dose one Hepatitis A dose 2 6 months later
12-15 months	Japanese encephalitis vaccine dose two Rabies dose one Rabies dose 2 after one week Rabies dose 3 after 3 weeks
	Diphtheria- Tetanus-acellular Pertussis-inactivated Polio MMR –measles mumps and Rubella
4-6 years	Diphtheria- Tetanus-acellular Pertussis-inactivated Polio
	MMR –measles mumps and Rubella
12-23 months Hepatitis A two doses given at least six months apart.	
Girls 12-13 Years Human papillomavirus (HPV) ( <a href="#">Cervarix</a> ) or Gardasil Three injections given over six months	
13-18 years Diphtheria + Tetanus + polio	
This table is provided for guidance only and space restrictions exclude regimes for Japanese encephalitis, Rabies, Varicella Influenza and Typhoid. Ask our staff for assistance * For Children at higher risk of TB ** Babies of Hepatitis B carriers need HBIG and hepatitis B vaccine at birth	